UNIVERSITY OF AGRICULTURAL SCIENCES, BENGALURU & INDIAN METEOROLOGICAL DEPARTMENT



GRAMIN KRISHI MAUSAM SEWA AMFU, OFRS, NAGANAHALLI, MYSURU - 570003



Date: 09-08-2024

AGRO-ADVISORY BULLETIN FOR MYSURU DISTRICT

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Past Weather Data

Parameter	06.08.2024	07.08.2024	08.08.2024	09.08.2024				
Rainfall (mm)	0	0	0.8	0				
Max. Temp. (°C)	30.3	29.2	27.2	28.7				
Min. Temp. (°C)	-	-	-	-				
Sky condition (Octas)	8	8	8	5				
Relative humidity (%) 0830 hours	81	77	79	78				
Relative humidity (%) 1730 hours	86	86	80	80				
Wind Speed (km/h)	6	6	6	8				
Wind Direction	230	230	230	230				

Weather forecast for the next five days (From 10-08-2024 to 14-08-2024)							
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024		
Rainfall (mm)	7	9	7	6	5		
Max. temp (°C)	29	30	31	31	31		
Min.Temp (°C)	16.8	17.3	17.8	18.2	17.5		
Sky condition (Octas)	8	8	8	8	8		
Relative humidity (%) 0830 hours	94	93	92	91	91		
Relative humidity (%) 1730 hours	62	62	60	56	59		
Wind Speed (kmph)	11	10	9	8	9		
Wind Direction	252	270	288	279	246		

Forecast Summary

As forecast received from IMD, cloudy sky with light to moderate rainfall may be expected from 10.08.2024 to 14.08.2024 in Mysuru district. The day temperature is expected to be 29-31°C & night temperature is expected 16.8-18.2°C. The relative humidity in the morning hours is expected to be 91-94% & afternoon relative humidity is expected to be in the range of 56-62%. Wind speed expected to be 8-11 km/hr.

Recommendations to the farmers:					
Crop	Pest/Disease	Damage symptoms	Control measures		
General Advisory					

• General recommendations for agricultural activities based on the given rainfall forecast:

- ✓ Postpone sowing if the soil moisture is insufficient due to light rainfall.
- ✓ Ensure proper soil moisture before sowing by irrigating if necessary.
- ✓ Carry out weeding during dry spells between rainfall events.
- ✓ Avoid weeding immediately after rain to prevent soil compaction.
- ✓ Limit irrigation due to forecasted light rainfall, but monitor soil moisture levels.
- ✓ Provide light irrigation if the rainfall is not sufficient to meet crop water requirements.
- ✓ Schedule spraying during dry periods, preferably in the early morning or late afternoon.
- ✓ Avoid spraying immediately before or after rainfall to ensure the effectiveness of the application.
- ✓ Plan harvesting during dry spells to prevent post-harvest losses due to moisture.
- ✓ Avoid harvesting when rainfall is forecasted to reduce the risk of crop damage.
- ✓ Ensure good drainage systems are in place to prevent waterlogging, especially in low-lying areas.
- ✓ Regularly inspect and clear drainage channels to allow excess water to flow out.
- ✓ Prune plants during dry days to minimize the risk of disease entry through fresh cuts.
- ✓ Avoid pruning when heavy rainfall is expected to reduce the risk of fungal infections.
- ✓ Apply fertilizers when light rainfall is forecasted to ensure better absorption by the soil.
- ✓ Avoid fertilizing just before heavy rainfall to prevent nutrient leaching.
- ✓ Treat seeds before sowing, especially if sowing is planned after light rainfall.
- ✓ Ensure treated seeds are dried properly before sowing to avoid fungal infections.

Crop	Stage	Advisory
Tomato	Ripening	Pest Management: Use spinosad. Monitor and manage fungal diseases.
Ginger, Turmeric	Rhizome	Ensure proper drainage. Use fungicides if necessary. Mulch
& Cabbage	development	to conserve soil moisture.
Pulses (Green		Use neem oil for pest control. Apply nitrogenous fertilizers
gram, Black gram,	Vegetative	if yellowing is observed.
Cowpea, Avare,		
Groundnut)	D1-	Annin in in it is a contract to the contract t
Banana	Bunch	Apply balanced fertilizers. Check for banana stem weevil and use control measures.
B 11 1B 1	development	
Paddy and Ragi	Transplanted	Ensure adequate irrigation and avoid water stagnation. Manual weeding or appropriate herbicides.
Coffee	Berry	Monitor for leaf rust and use recommended fungicides.
	development	Ensure adequate shade.
Maize, Brinjal,		Monitor for aphids, jassids, and whiteflies. Use appropriate
Sunflower, Chilli	Vegetative	insecticides.
and Cotton		
Beans	Flowering	Monitor for aphids and bean fly. Use neem-based products.
	Flowering	Ensure regular irrigation.
Coconut &		Use neem oil for rugose whitefly. Remove affected leaves
Arecanut	-	for caterpillar. Use pheromone traps for rhinoceros beetle.

Maize Fall army worm	Vegetative stage	Use chlorantraniliprole or Emamectin benzoate for fall armyworm. Regular monitoring.
Tobacco	Harvesting	Ensure proper curing conditions to prevent mold and rot. Avoid harvesting during rainy periods. Handle leaves gently to prevent damage.
Tomato Fruit borer	Fruiting	 Caterpillar bore the flower buds and fruits. Infested flower buds with hole and drops off, fruit with a hole, water enter through the hole leads to rotting. Trap crop: For every 25 rows of tomato grow one row of marigold cultivar African tall. The marigold seedlings about 35-40 days old. If borer problems exceeds10 per cent spray 4 per cent. NSKE or 100 LE, Ha. NPV. If infestation in severe form spray 1.0 g. Methomil 40 SP. in a lit. water
Coconut black headed caterpillar	 On conpalm the Method: A management of the position. To breadth add quantity of we insecticide minut after a mongravid, Caution: No. 	e and burn the severly affected fronds. Inmunity basis feed the Manocrotophos 36 SL. to the rough root. Ineter away from trunk, dig out and select brown cil thickness size root. Cut the root in a slanting the polythene bag (size of 15 cm. length 4 cm. 7.5 to 10 ml. Monocrotophos 36 SL. with equal rater, introduce and immerse cut end of the root in xture and tie the bag with thread. In absorb the chemical within a period of 24 hours, if r 48 hours select another root to feed the chemical. The after chemial treatment release larval parasites: Goniozus@ 10 - 12 /palm. In to harvest tender coconuts/matured coconuts of the date of chemical treatment.
Coconut	Rhinoceros beetle	Remove the beetle from infected part and fill 2 % Quinolphos or 5 % Melathion in sand @ 1:1 ratio.
Sericulture	Provide good	al temperature and humidity in silkworm rearing houses. ventilation. Use disinfectants to control diseases. Ensure are clean and disease-free.
Poultry	and humidity.	Provide clean drinking water. Vaccinate against common the poultry shed dry and clean to prevent disease outbreaks.
Livestock	clean drinking v	nd dry housing to prevent hoof diseases. Provide adequate vater. Supplement diet with minerals and vitamins. Regularly al parasites and treat accordingly.

Block level weather forecast (From 10-08-2024 to 14-08-2024)

H.D. Kote						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	6.5	18.4	8.2	11.8	9.1	
Max. temp (°C)	27.3	28.4	28.5	29.1	26.6	
Min.Temp (°C)	19	19.6	19.9	20.3	19.9	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	96	95	95	92	92	
Relative humidity (%) 1730 hours	62	59	61	56	59	
Wind Speed (kmph)	12	11	9	10	9	
Wind Direction	248	252	249	257	248	

Hunsuru						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	16.6	23.1	16.1	16.1	13.5	
Max. temp (°C)	28.3	28.8	29.3	29.2	27.1	
Min.Temp (°C)	18.8	19.2	19.6	19.8	19.7	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	95	94	93	90	92	
Relative humidity (%) 1730 hours	60	60	60	54	59	
Wind Speed (kmph)	11	11	9	10	9	
Wind Direction	248	288	289	283	257	

K.R. Nagara						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	4.2	15.2	0	2.8	16.7	
Max. temp (°C)	24	27.1	27.2	27.6	28.1	
Min.Temp (°C)	17.9	18	17.8	17.6	18.8	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	93	91	91	94	92	
Relative humidity (%) 1730 hours	74	62	62	55	54	
Wind Speed (kmph)	10	14	13	12	11	
Wind Direction	249	252	257	249	283	

Mysuru						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	22.5	55.3	37.6	23	17.7	
Max. temp (°C)	27.9	27.7	28.2	28.4	26.2	
Min.Temp (°C)	18.2	18.7	18.9	19.3	19.1	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	94	92	93	90	90	
Relative humidity (%) 1730 hours	55	59	58	56	57	
Wind Speed (kmph)	13	12	10	10	9	
Wind Direction	248	283	250	283	257	

Nanjanagudu						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	8.5	38	18	20.5	8.9	
Max. temp (°C)	26.8	27	27.1	27.2	24.8	
Min.Temp (°C)	17.5	18.1	18.2	18.8	18.5	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	92	92	92	89	89	
Relative humidity (%) 1730 hours	53	55	55	54	57	
Wind Speed (kmph)	14	13	11	10	9	
Wind Direction	249	252	248	257	249	

Piriapatna						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	10.4	7.1	12.8	15.3	10.6	
Max. temp (°C)	28.5	29.4	30.2	29.6	27.6	
Min.Temp (°C)	18.7	19.2	19.5	19.7	19.7	
Sky condition (Octas)	8	8	8	8	8	
Relative humidity (%) 0830 hours	96	93	93	91	93	
Relative humidity (%) 1730 hours	62	60	60	53	58	
Wind Speed (kmph)	10	11	10	11	9	
Wind Direction	249	288	283	288	257	

T. Narasipura						
Parameter	10.08.2024	11.08.2024	12.08.2024	13.08.2024	14.08.2024	
Rainfall (mm)	9.3	30	13.1	14.8	2.7	
Max. temp (°C)	27.8	28.1	28	27.5	25.6	
Min.Temp (°C)	18.3	18.9	19	19.6	19.5	
Sky condition (Octas)	7	8	8	8	8	
Relative humidity (%) 0830 hours	91	88	91	87	86	
Relative humidity (%) 1730 hours	51	52	51	53	54	
Wind Speed (kmph)	14	11	11	10	9	
Wind Direction	252	261	250	257	249	

- Download "**DAMINI**" app to get early warning on lightening and take precautions based on the alert given by the application.
- ➤ Kindly download "MAUSAM" APP for location specific forecast & warning & "MEGHDOOT" APP for Agromet advisory
- This information is available in the website: mausam.imd.gov.in

For any information farmers can contact **Dr. C. Ramachandra**, Senior Farm Superintendent/ **Dr. Sumanth Kumar.G.V**, Technical officer over phone No. 0821-259126/9535345814.

AMFU of IMD, Naganahalli, Mysuru

वास्तविकवर्षातथाविस्तारितअवधिपूर्वानुमान Realized Rainfall and Extended Range Forecast (वर्षाऔरतापमान)

(Rainfall and Temperature)

Realized Rainfall (18thto 31stJuly,2024)







- Normal or above normal rainfall occurred in both the weeks over Uttarakhand, West Rajasthan, Madhya Pradesh, Chhattisgarh, Odisha, Gujarat Region, Maharashtra, Telangana, Coastal Andhra Pradesh, Karnataka, Kerala & Mahe and Lakshadweep.
- Normal or above normal rainfall occurred in either of the two weeks over Jammu & Kashmir and Ladakh (UTs), Himachal Pradesh, West Uttar Pradesh, Jharkhand, Gangetic West Bengal, Saurashtra & Kutch, East Rajasthan, Rayalaseema and Tamil Nadu Puducherry & Karaikal.
- Below Normal rainfall/no rain occurred in both the weeks over Punjab, Haryana Chandigarh & Delhi, East Uttar Pradesh, Bihar, Sub-Himalayan West Bengal & Sikkim, Assam & Meghalaya, Arunachal Pradesh, Nagaland Manipur Mizoram & Tripura (NMMT) and Andaman & Nicobar Islands.

Extended Range Forecast System


Rainfall forecast maps for the next 2 weeks (IC- 31stJuly, 2024) (02ndto 15thAugust, 2024)






- Week 1 (02.08.2024 to 08.08.2024): Rainfall is likely to be above normal over most parts of the country.
- Week 2 (09.08.2024 to 15.08.2024):Rainfall is likely to be above normal overNorthwest India and Indo Gangetic Plains. However, it is likely to be below normal over Central India, Northeast India and along West coast.

Maximum and Minimum temperature anomaly ($^{\circ}$ C) forecast for the next 2 weeks (IC- 31^{st} July, 2024) $(02^{nd}to\ 15^{th}August,\ 2024)$



Maximum Temperature (Tmax)

- Week 1 (02.08.2024 to 08.08.2024): Maximum temperature is likely to be normal to below normal over most parts of the country.
- Week 2 (09.08.2024 to 15.08.2024): Maximum temperature is likely to above normal over South India and some parts of Central India. It is likely to be below normal over Northwest India, East India and Northeast India.



Minimum Temperature (Tmin)

- Week 1 (02.08.2024 to 08.08.2024): Minimum temperature likely to be slightly above normal over parts of Northwest India, Central India, East India and Karnataka.
- Week 2 (09.08.2024 to 15.08.2024): Minimum temperature likely to be slightly above normal over parts of Maharashtra, Madhya Pradesh, Chhattisgarh, Gujarat, Himachal Pradesh and Uttarakhand. It is likely to be above normal over parts of Karnataka.